



# Asymptotic models for the multiple electromagnetic wave scattering problem by small obstacles

Justine Labat, Victor Péron, Sébastien Tordeux

## ► To cite this version:

Justine Labat, Victor Péron, Sébastien Tordeux. Asymptotic models for the multiple electromagnetic wave scattering problem by small obstacles. Journées Ondes Sud-Ouest (JOSO), Mar 2019, Le Barp, France. hal-02071888

**HAL Id: hal-02071888**

**<https://inria.hal.science/hal-02071888>**

Submitted on 19 Mar 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Asymptotic models for the multiple electromagnetic wave scattering problem by small obstacles

Justine Labat, Victor Péron, Sébastien Tordeux

PhD student in Applied Mathematics


EPC Magique 3D — UPPA-E2S, INRIA Bordeaux Sud-Ouest, LMAP UMR CNRS 5142

Journées Ondes Sud-Ouest

Le Barp, March 13, 2019




# 3D Scattering problem by small obstacles



Asymptotic assumption :  $\delta \ll \lambda$

# Model problem



- Time-harmonic domain
- Homogeneous & isotropic medium  $\Omega_\delta$
- Perfect conductors of characteristic length  $\delta$

## Time-harmonic Maxwell equations

$$\begin{cases} \mathbf{curl} \mathbf{E}_\delta - i\kappa \mathbf{H}_\delta = 0 & \text{in } \Omega_\delta \\ \mathbf{curl} \mathbf{H}_\delta + i\kappa \mathbf{E}_\delta = 0 & \text{in } \Omega_\delta \end{cases}$$

$$\text{with } \kappa^2 = \omega^2 \mu \left( \epsilon + \frac{i\sigma}{\omega} \right), \Im(\kappa) \geq 0$$

## Boundary condition

$$\mathbf{n} \times \mathbf{E}_\delta = -\mathbf{n} \times \mathbf{E}^{\text{inc}} \quad \text{on } \partial\Omega_\delta$$

## Silver-Müller radiation condition

$$r(\mathbf{H}_\delta \times \hat{\mathbf{x}} - \mathbf{E}_\delta) \xrightarrow[r \rightarrow \infty]{} 0 \quad \text{unif. in } \hat{\mathbf{x}} = \frac{\mathbf{x}}{r}$$

## Mathematical well-posedness

For any  $\mathbf{E}^{\text{inc}} \in \mathbf{H}_{\text{loc}}(\mathbf{curl}, \Omega_\delta)$  there exist a unique solution  $(\mathbf{E}_\delta, \mathbf{H}_\delta) \in \mathbf{H}_{\text{loc}}(\mathbf{curl}, \Omega_\delta)^2$  to the exterior Maxwell problem.

# Applications

- Medical imaging
- Civil engineering
- Nuclear industry
- Atmospheric science
- Astronomy
- Climatology
- ...

## Inverse problem ?

- Locate heterogeneities, cracks, ...
- Characterize them




Figure: Small tumor detection<sup>1</sup>



Figure: Non-destructive testing<sup>2</sup>

<sup>1</sup> <https://medcitynews.com/wp-content/uploads/2017/01/GettyImages-94456546-600x450.jpg>

<sup>2</sup> <https://tri-intl.com/wp-content/uploads/2017/07/NonDestructiveTesting.jpg>

## Asymptotic models

- ✗ Restricted to small obstacles
- ✓ Low computational cost
- ✓ Meshless method

# Workflow

## Asymptotic models

- ✗ Restricted to small obstacles
- ✓ Low computational cost
- ✓ Meshless method

Multiple scattering



## Foldy-Lax model

- ✓ Interactions taken into account
- ✓ Low computational cost
- ✓ Meshless method

Superposition  
principle



## Born approximation

- ✗ No interaction between the obstacles
- ✓ Low computational cost
- ✓ Meshless method

# Workflow

## Asymptotic models

- ✗ Restricted to small obstacles
- ✓ Low computational cost
- ✓ Meshless method

Multiple scattering



## Foldy-Lax model

- ✓ Interactions taken into account
- ✓ Low computational cost
- ✓ Meshless method

Superposition  
principle



## Born approximation

- ✗ No interaction between the obstacles
- ✓ Low computational cost
- ✓ Meshless method

### (Non-exhaustive) alternatives

- Boundary integral equations
  - ▶ Boundary element methods
  - ▶ Spectral-based methods
- Enrichment of approximation spaces
  - ▶ Trefftz-based methods
  - ▶ Extended FEM
  - ▶ Partition of Unity method



# Workflow

## Asymptotic models

- ✗ Restricted to small obstacles
- ✓ Low computational cost
- ✓ Meshless method

Multiple scattering



## Foldy-Lax model

- ✓ Interactions taken into account
- ✓ Low computational cost
- ✓ Meshless method

Superposition  
principle



## Born approximation

- ✗ No interaction between the obstacles
- ✓ Low computational cost
- ✓ Meshless method

### (Non-exhaustive) alternatives

- Boundary integral equations
  - ▶ Boundary element methods
  - ✓ Spectral-based methods
- Enrichment of approximation spaces
  - ▶ Trefftz-based methods
  - ▶ Extended FEM
  - ▶ Partition of Unity method

# Some references

- Historic references
  - ▶ Rayleigh (1884), Foldy (1945), Lax (1951)
- **Small defect** theory
  - ▶ Il'in (1992), Maz'ya-Nazarov-Plamenevskii (2000)
- Acoustic obstacle
  - ▶ Ammari-Kang (2003), Ramm (2005), Claeys (2008)
- Time-dependent domain
  - ▶ Mattesi (2014), Korikov (2015), Marmorat (2015)
- **Electromagnetic** obstacle
  - ▶ Vogelius-Volkov (2000), Ammari-Vogelius-Volkov (2001), Korikov-Plamenevskii (2017)
- **Foldy** theory
  - ▶ Martin (2004), Cassier-Hazard (2013), Bendali-Cocquet-Tordeux (2014), Challa-Hu-Sini (2014)
- High-order **spectral** algorithms
  - ▶ Xu (1995), Ganesh-Hawkins (2009), Barucq-Chabassier-Pham-Tordeux (2017)
- Inverse problem
  - ▶ Volkov (2001), Ammari-Kang (2004), Challa-Sini (2012)

# Some references

- Historic references
  - ▶ Rayleigh (1884), Foldy (1945), Lax (1951)
- **Small defect** theory
  - ▶ Il'in (1992), Maz'ya-Nazarov-Plamenevskii (2000)
- Acoustic obstacle
  - ▶ Ammari-Kang (2003), Ramm (2005), Claeys (2008)
- Time-dependent domain
  - ▶ Mattesi (2014), Korikov (2015), Marmorat (2015)
- **Electromagnetic** obstacle
  - ▶ Vogelius-Volkov (2000), Ammari-Vogelius-Volkov (2001), Korikov-Plamenevskii (2017)
- **Foldy** theory
  - ▶ Martin (2004), Cassier-Hazard (2013), Bendali-Cocquet-Tordeux (2014), Challa-Hu-Sini (2014)
- High-order **spectral** algorithms
  - ▶ Xu (1995), Ganesh-Hawkins (2009), Barucq-Chabassier-Pham-Tordeux (2017)
- Inverse problem
  - ▶ Volkov (2001), Ammari-Kang (2004), Challa-Sini (2012)

# Outline

## 1. Asymptotic models

- Single-scattering
- Application: Born approximation
- Multiple-scattering: Foldy-Lax model
- Numerical results

## 2. Spectral method: Spherical case

- Discretization
- Numerical convergence
- Comparison with asymptotic models
- Comparison with finite element solutions

## 3. Conclusions and perspectives

# Outline

## 1. Asymptotic models

- Single-scattering
- Application: Born approximation
- Multiple-scattering: Foldy-Lax model
- Numerical results

## 2. Spectral method: Spherical case



- Discretization
- Numerical convergence
- Comparison with asymptotic models
- Comparison with finite element solutions

## 3. Conclusions and perspectives

# Approximation of solution to single scattering

## Method of **matched asymptotic expansions**



- Domain decomposition
- Local approximations
- Matching procedure



# Approximation of solution to single scattering

## Method of **matched asymptotic expansions**

- Domain decomposition
- **Local approximations**
- Matching procedure





## Asymptotic expansions

- **Far field** expansion in  $\Omega^* = \mathbb{R}^3 \setminus \{0\}$
- **Near field** expansion in  $\hat{\Omega} = \mathbb{R}^3 \setminus \overline{\mathcal{B}(0, 1)}$

# Approximation of solution to single scattering

## Method of **matched asymptotic expansions**

- Domain decomposition
- Local approximations
- **Matching procedure**



## Asymptotic expansions

- **Far field** expansion in  $\Omega^* = \mathbb{R}^3 \setminus \{0\}$
- **Near field** expansion in  $\hat{\Omega} = \mathbb{R}^3 \setminus \overline{\mathcal{B}(0, 1)}$

related by an asymptotic process



# Asymptotic expansions

## Near field expansion

$$\mathbf{E}_\delta(\delta\mathbf{X}) \approx \widehat{\mathbf{E}}_0(\mathbf{X}) + \delta\widehat{\mathbf{E}}_1(\mathbf{X}) + \delta^2\widehat{\mathbf{E}}_2(\mathbf{X}) + \dots$$

$$\mathbf{H}_\delta(\delta\mathbf{X}) \approx \widehat{\mathbf{H}}_0(\mathbf{X}) + \delta\widehat{\mathbf{H}}_1(\mathbf{X}) + \delta^2\widehat{\mathbf{H}}_2(\mathbf{X}) + \dots$$

$\mathbf{X} = \frac{\mathbf{x}}{\delta}$ : fast variable

## Far field expansion

$$\mathbf{E}_\delta(\mathbf{x}) \approx \delta^3\widetilde{\mathbf{E}}_3(\mathbf{x}) + \delta^4\widetilde{\mathbf{E}}_4(\mathbf{x}) + \delta^5\widetilde{\mathbf{E}}_5(\mathbf{x}) + \dots$$

$$\mathbf{H}_\delta(\mathbf{x}) \approx \delta^3\widetilde{\mathbf{H}}_3(\mathbf{x}) + \delta^4\widetilde{\mathbf{H}}_4(\mathbf{x}) + \delta^5\widetilde{\mathbf{H}}_5(\mathbf{x}) + \dots$$

- For an obstacle of arbitrary shape

Numerical solution of elementary problems **independent of  $\delta$**

- ▶ Near-field: **quasi-static** problems
- ▶ Far-field: **time-harmonic** problems + equivalent multipole distributions

- For a **spherical** obstacle

- ▶ Analytical solutions of elementary problems
- ▶ Equivalent **multipole distributions**

# Near-field approximation: Spherical case

Close to the obstacle : **quasi-static** approximation

Superposition of electric fields generated by **multipole distributions**:


$$\hat{\mathbf{E}}_p = \sum_{\substack{n=1 \\ |n-p| \text{ odd}}}^{p+1} \mathcal{E}_{n,\text{elec}}^{\text{stat}}[\mathbf{u}_{1,n}^{p,E}, \dots, \mathbf{u}_{n,n}^{p,E}] + \sum_{\substack{n=1 \\ |n-p| \text{ even}}}^{p+1} \mathcal{E}_{n,\text{mag}}^{\text{stat}}[\mathbf{u}_{1,n}^{p,H}, \dots, \mathbf{u}_{n,n}^{p,H}] \quad p = 0, 1, 2, \dots$$

Dipole ( $n = 1$ )

Quadrupole ( $n = 2$ )

$\mathcal{E}_n^{\text{stat}}[\mathbf{d}_1, \dots, \mathbf{d}_n]$  electric fields

Charges



Currents



- generated by **charges** OR **currents**
- electric/magnetic  **$2^n$ -point charges**
  - ▶ electric charge:  $\mathcal{E} = -\nabla \mathcal{V}$
  - ▶ magnetic current:  $\mathcal{E} = -\text{curl } \mathcal{A}$
- of **moments** ( $\mathbf{d}_1, \dots, \mathbf{d}_n$ )
- defined by an **asymptotic** process

# Near-field approximation: Spherical case

Close to the obstacle : **quasi-static** approximation

Superposition of electric fields generated by **multipole distributions**:

$$\hat{\mathbf{E}}_p = \sum_{\substack{n=1 \\ |n-p| \text{ odd}}}^{p+1} \boldsymbol{\varepsilon}_{n,\text{elec}}^{\text{stat}}[\mathbf{u}_{1,n}^{p,E}, \dots, \mathbf{u}_{n,n}^{p,E}] + \sum_{\substack{n=1 \\ |n-p| \text{ even}}}^{p+1} \boldsymbol{\varepsilon}_{n,\text{mag}}^{\text{stat}}[\mathbf{u}_{1,n}^{p,H}, \dots, \mathbf{u}_{n,n}^{p,H}] \quad p = 0, 1, 2, \dots$$

	Order	Dipole		Quadrupole		Octupole	
		Electric	Magnetic	Electric	Magnetic	Electric	Magnetic
Near field	0	●	●				
	1	●	●	●	●		
	2	●	●	●	●	●	●

● Electric field

● Magnetic field


# Far-field approximation

Far from the obstacle: **time-harmonic** approximation


Superposition of electric fields generated by **multipole distributions**:

$$\tilde{\mathbf{E}}_{p+3} = \sum_{n=1}^{p+1} \boldsymbol{\varepsilon}_{n,\text{elec}}[\mathbf{v}_{1,n}^{p,\text{E}}, \dots, \mathbf{v}_{n,n}^{p,\text{E}}] + \boldsymbol{\varepsilon}_{n,\text{mag}}[\mathbf{v}_{1,n}^{p,\text{H}}, \dots, \mathbf{v}_{n,n}^{p,\text{H}}] \quad p = 0, 1, 2, \dots$$

Dipole ( $n = 1$ )



Quadrupole ( $n = 2$ )



$\boldsymbol{\varepsilon}_n[\mathbf{d}_1, \dots, \mathbf{d}_n]$  electric fields

- generated by charges **AND** currents
  - related by **charge conservation principle**
- electric/magnetic  $2^n$ -point charges
  - electric multipole:  $\boldsymbol{\varepsilon} = -\nabla \mathcal{V}_\text{E} + i\kappa \mathcal{A}_\text{E}$
  - magnetic multipole:  $\boldsymbol{\varepsilon} = -\text{curl } \mathcal{A}_\text{H}$
- of moments ( $\mathbf{d}_1, \dots, \mathbf{d}_n$ )
- defined by an asymptotic process

$$\text{div } \mathcal{J} - i\omega \varrho = 0$$

# Far-field approximation

Far from the obstacle: **time-harmonic** approximation

$$\tilde{\mathbf{E}}_{p+3} = \sum_{n=1}^{p+1} \boldsymbol{\varepsilon}_{n,\text{elec}}[\mathbf{v}_{1,n}^{p,E}, \dots, \mathbf{v}_{n,n}^{p,E}] + \boldsymbol{\varepsilon}_{n,\text{mag}}[\mathbf{v}_{1,n}^{p,H}, \dots, \mathbf{v}_{n,n}^{p,H}]$$

$$\tilde{\mathbf{H}}_{p+3} = \sum_{n=1}^{p+1} \boldsymbol{\varkappa}_{n,\text{mag}}[\mathbf{v}_{1,n}^{p,H}, \dots, \mathbf{v}_{n,n}^{p,H}] + \boldsymbol{\varkappa}_{n,\text{elec}}[\mathbf{v}_{1,n}^{p,E}, \dots, \mathbf{v}_{n,n}^{p,E}]$$

	Order	Dipole				Quadrupole				Octupole			
		Electric		Magnetic		Electric		Magnetic		Electric		Magnetic	
Far field	3	●	●	●	●								
	4	●	●	●	●	●	●	●	●				
	5	●	●	●	●	●	●	●	●	●	●	●	●

● Electric field

● Magnetic field

# Far-field approximation

Far from the obstacle: **time-harmonic** approximation

$$\tilde{\mathbf{E}}_{p+3} = \sum_{n=1}^{p+1} \boldsymbol{\varepsilon}_{n,\text{elec}}[\mathbf{v}_{1,n}^{p,E}, \dots, \mathbf{v}_{n,n}^{p,E}] + \boldsymbol{\varepsilon}_{n,\text{mag}}[\mathbf{v}_{1,n}^{p,H}, \dots, \mathbf{v}_{n,n}^{p,H}]$$

$$\tilde{\mathbf{H}}_{p+3} = \sum_{n=1}^{p+1} \boldsymbol{\varkappa}_{n,\text{mag}}[\mathbf{v}_{1,n}^{p,H}, \dots, \mathbf{v}_{n,n}^{p,H}] + \boldsymbol{\varkappa}_{n,\text{elec}}[\mathbf{v}_{1,n}^{p,E}, \dots, \mathbf{v}_{n,n}^{p,E}]$$

- Spherical** case

	Order	Dipole		Quadrupole		Octupole	
		Electric	Magnetic	Electric	Magnetic	Electric	Magnetic
Far field	3	•	•	•	•		
	4						
	5	•	•	•	•	•	•

- Collected dipolar** approximation

	Order	Dipole		Quadrupole		Octupole	
		Electric	Magnetic	Electric	Magnetic	Electric	Magnetic
Far field	3	•	•	•	•		
	4						
	5	•	•	•	•		

# Born approximation

- The electromagnetic fields are approximated by the **superposition principle**

$$\mathbf{E}_\delta(\mathbf{x}) \approx \sum_{k=1}^{N_{\text{obs}}} \mathbf{E}_{\delta,k}(\mathbf{x}) \quad \mathbf{H}_\delta(\mathbf{x}) \approx \sum_{k=1}^{N_{\text{obs}}} \mathbf{H}_{\delta,k}(\mathbf{x})$$

- Each obstacle is modeled as a **dipolar source** around  $c_k$

$$\mathbf{E}_{\delta,k}(\mathbf{x}) = \boldsymbol{\varepsilon}_{1,\text{elec}}[\mathbf{d}_{\delta,k}^{\text{E}}](\mathbf{x} - c_k) + \boldsymbol{\varepsilon}_{1,\text{mag}}[\mathbf{d}_{\delta,k}^{\text{H}}](\mathbf{x} - c_k)$$

For **perfectly conducting spheres**:

- Approximation of order 3

$$\mathbf{d}_{\delta,k}^{\text{E}} = 4\pi\delta^3 \mathbf{E}^{\text{inc}}(c_k) \quad \mathbf{d}_{\delta,k}^{\text{H}} = -2\pi\delta^3 \mathbf{H}^{\text{inc}}(c_k)$$

- Collected dipolar approximation

$$\mathbf{d}_{\delta,k}^{\text{E}} = 4\pi\delta^3 \left(1 + \frac{3(\kappa\delta)^2}{10}\right) \mathbf{E}^{\text{inc}}(c_k) \quad \mathbf{d}_{\delta,k}^{\text{H}} = -2\pi\delta^3 \left(1 - \frac{3(\kappa\delta)^2}{5}\right) \mathbf{H}^{\text{inc}}(c_k)$$

# Foldy-Lax model

- The electromagnetic fields are approximated by the superposition principle

$$\mathbf{E}_\delta(\mathbf{x}) \approx \sum_{k=1}^{N_{\text{obs}}} \mathbf{E}_{\delta,k}(\mathbf{x}) \quad \mathbf{H}_\delta(\mathbf{x}) \approx \sum_{k=1}^{N_{\text{obs}}} \mathbf{H}_{\delta,k}(\mathbf{x})$$

- Each obstacle is modeled as a dipolar source around  $c_k$

$$\mathbf{E}_{\delta,k}(\mathbf{x}) = \boldsymbol{\varepsilon}_{1,\text{elec}}[\mathbf{d}_{\delta,k}^{\text{E}}](\mathbf{x} - c_k) + \boldsymbol{\varepsilon}_{1,\text{mag}}[\mathbf{d}_{\delta,k}^{\text{H}}](\mathbf{x} - c_k)$$

For **perfectly conducting** spheres:

- Approximation of order 3

$$\mathbf{d}_{\delta,k}^{\text{E}} = 4\pi\delta^3 \left( \mathbf{E}^{\text{inc}}(c_k) + \sum_{\substack{\ell=1 \\ \ell \neq k}}^{N_{\text{obs}}} \mathbf{E}_{\delta,\ell}(c_k) \right) \quad \mathbf{d}_{\delta,k}^{\text{H}} = -2\pi\delta^3 \left( \mathbf{H}^{\text{inc}}(c_k) + \sum_{\substack{\ell=1 \\ \ell \neq k}}^{N_{\text{obs}}} \mathbf{H}_{\delta,\ell}(c_k) \right)$$

- Collected dipolar approximation

$$\mathbf{d}_{\delta,k}^{\text{E}} = 4\pi\delta^3 \left( 1 + \frac{3(\kappa\delta)^2}{10} \right) \left( \mathbf{E}^{\text{inc}}(c_k) + \sum_{\substack{\ell=1 \\ \ell \neq k}}^{N_{\text{obs}}} \mathbf{E}_{\delta,\ell}(c_k) \right) \quad \mathbf{d}_{\delta,k}^{\text{H}} = -2\pi\delta^3 \left( 1 - \frac{3(\kappa\delta)^2}{5} \right) \left( \mathbf{H}^{\text{inc}}(c_k) + \sum_{\substack{\ell=1 \\ \ell \neq k}}^{N_{\text{obs}}} \mathbf{H}_{\delta,\ell}(c_k) \right)$$



# Foldy-Lax model

- The electromagnetic fields are approximated by the superposition principle

$$\mathbf{E}_\delta(\mathbf{x}) \approx \sum_{k=1}^{N_{\text{obs}}} \mathbf{E}_{\delta,k}(\mathbf{x}) \quad \mathbf{H}_\delta(\mathbf{x}) \approx \sum_{k=1}^{N_{\text{obs}}} \mathbf{H}_{\delta,k}(\mathbf{x})$$

- Each obstacle is modeled as a dipolar source around  $c_k$

$$\mathbf{E}_{\delta,k}(\mathbf{x}) = \boldsymbol{\varepsilon}_{1,\text{elec}}[\mathbf{d}_{\delta,k}^{\text{E}}](\mathbf{x} - c_k) + \boldsymbol{\varepsilon}_{1,\text{mag}}[\mathbf{d}_{\delta,k}^{\text{H}}](\mathbf{x} - c_k)$$

- Vectorial formulation** (3-order approximation)

Find  $\mathbf{d} = ((\mathbf{d}_1^{\text{E}}), \dots, (\mathbf{d}_{N_{\text{obs}}}^{\text{E}}), (\mathbf{d}_1^{\text{H}}), \dots, (\mathbf{d}_{N_{\text{obs}}}^{\text{H}}))^{\top} \in \mathbb{C}^{6N_{\text{obs}}}$  such that

$$\mathbf{d} - \delta^3 \mathbb{A} \mathbf{d} = \delta^3 \mathbf{f}$$

$$\text{with } \mathbf{f} = \begin{pmatrix} 4\pi \mathbf{E}^{\text{inc}}(c_1) \\ \vdots \\ 4\pi \mathbf{E}^{\text{inc}}(c_{N_{\text{obs}}}) \\ -2\pi \mathbf{H}^{\text{inc}}(c_1) \\ \vdots \\ -2\pi \mathbf{H}^{\text{inc}}(c_{N_{\text{obs}}}) \end{pmatrix}$$

and  $\mathbb{A}$  the “interaction” matrix




Figure: Skeleton of  $\mathbb{A}$  for  $N_{\text{obs}} = 4$

# Foldy-Lax approximations

- Approximation of order 3

$$\mathbf{d} - \delta^3 \mathbb{A} \mathbf{d} = \delta^3 \mathbf{f}$$

- Collected dipolar approximation

$$\mathbb{D}^{-1} \mathbf{d} - \delta^3 \mathbb{A} \mathbf{d} = \delta^3 \mathbf{f}$$

where

$$\mathbb{D} = \text{diag}(\underbrace{\alpha, \dots, \alpha}_{3N_{\text{obs}}}, \underbrace{\beta, \dots, \beta}_{3N_{\text{obs}}}) \quad \alpha = 1 + \frac{3(\kappa\delta)^2}{10} \quad \beta = 1 - \frac{6(\kappa\delta)^2}{10}$$

- Modified dipolar approximation

$$\tilde{\mathbb{D}}^{-1} \mathbf{d} - \delta^3 \mathbb{A} \mathbf{d} = \delta^3 \mathbf{f}$$

where

$$\tilde{\mathbb{D}} = \text{diag}(\underbrace{\tilde{\alpha}, \dots, \tilde{\alpha}}_{3N_{\text{obs}}}, \underbrace{\tilde{\beta}, \dots, \tilde{\beta}}_{3N_{\text{obs}}}) \quad \tilde{\alpha} = \frac{3i}{2(\kappa\delta)^3} \frac{j_1(\kappa\delta)}{h_1^{(1)}(\kappa\delta)} \quad \tilde{\beta} = -\frac{3i}{(\kappa\delta)^3} \frac{j_1(\kappa\delta) + \kappa\delta j_1'(\kappa\delta)}{h_1^{(1)}(\kappa\delta) + \kappa\delta h_1^{(1)'}(\kappa\delta)}$$

# Foldy-Lax approximations

- Approximation of order 3

$$\mathbf{d} - \delta^3 \mathbb{A} \mathbf{d} = \delta^3 \mathbf{f}$$

- Collected dipolar approximation

$$\mathbb{D}^{-1} \mathbf{d} - \delta^3 \mathbb{A} \mathbf{d} = \delta^3 \mathbf{f}$$

where

$$\mathbb{D} = \text{diag}(\underbrace{\alpha, \dots, \alpha}_{3N_{\text{obs}}}, \underbrace{\beta, \dots, \beta}_{3N_{\text{obs}}}) \quad \alpha = 1 + \frac{3(\kappa\delta)^2}{10} \quad \beta = 1 - \frac{6(\kappa\delta)^2}{10}$$

- **Modified** dipolar approximation


$$\tilde{\mathbb{D}}^{-1} \mathbf{d} - \delta^3 \mathbb{A} \mathbf{d} = \delta^3 \mathbf{f}$$

where


$$\tilde{\mathbb{D}} = \text{diag}(\underbrace{\tilde{\alpha}, \dots, \tilde{\alpha}}_{3N_{\text{obs}}}, \underbrace{\tilde{\beta}, \dots, \tilde{\beta}}_{3N_{\text{obs}}}) \quad \tilde{\alpha} = \frac{3i}{2(\kappa\delta)^3} \frac{j_1(\kappa\delta)}{h_1^{(1)}(\kappa\delta)} \quad \tilde{\beta} = -\frac{3i}{(\kappa\delta)^3} \frac{j_1(\kappa\delta) + \kappa\delta j_1'(\kappa\delta)}{h_1^{(1)}(\kappa\delta) + \kappa\delta h_1^{(1)'}(\kappa\delta)}$$

## Numerical validation: Spherical case

- Incident field: Electromagnetic plane wave of wavelength  $\lambda = 5$  along  $z$ -axis polarized by  $\mathbf{e}_x$




(a) Electric field ( $x$ -component)




(b) Magnetic field ( $y$ -component)


- Reference solution: “Spectral” solution truncated at the order 10



(c) One obstacle



(d) Aligned obstacles



(e) Real 3D-case



# Single-scattering: Validation of asymptotic expansions

- Near-field** approximations


Order 0 :  $\|\mathbf{E}_\delta - \widehat{\mathbf{E}}_0(\frac{\cdot}{\delta})\|_{\mathbf{L}^2(\Omega_\delta^{2\delta})} / \|\mathbf{E}_\delta\|_{\mathbf{L}^2(\Omega_\delta^{2\delta})}$

Order 1 :  $\|\mathbf{E}_\delta - \widehat{\mathbf{E}}_0(\frac{\cdot}{\delta}) - \delta \widehat{\mathbf{E}}_1(\frac{\cdot}{\delta})\|_{\mathbf{L}^2(\Omega_\delta^{2\delta})} / \|\mathbf{E}_\delta\|_{\mathbf{L}^2(\Omega_\delta^{2\delta})}$

Order 2 :  $\|\mathbf{E}_\delta - \widehat{\mathbf{E}}_0(\frac{\cdot}{\delta}) - \delta \widehat{\mathbf{E}}_1(\frac{\cdot}{\delta}) - \delta^2 \widehat{\mathbf{E}}_2(\frac{\cdot}{\delta})\|_{\mathbf{L}^2(\Omega_\delta^{2\delta})} / \|\mathbf{E}_\delta\|_{\mathbf{L}^2(\Omega_\delta^{2\delta})}$



(f) Electric field





(g) Magnetic Field

# Single-scattering: Validation of asymptotic expansions


- Far-field approximations

Order 3 :  $\|\mathbf{E}_\delta - \delta^3 \tilde{\mathbf{E}}_3\|_{\mathbf{L}^2(\Omega_\lambda^{2\lambda})}$

Order 5 :  $\|\mathbf{E}_\delta - \delta^3 \tilde{\mathbf{E}}_3 - \delta^5 \tilde{\mathbf{E}}_5\|_{\mathbf{L}^2(\Omega_\lambda^{2\lambda})}$




(l) Electric field




(m) Magnetic field

# Multiple-scattering: Validation of Foldy-Lax model


- Order 3: **Born** and **Foldy**
- Collected dipole improvement: **Collected Born** and **Collected Foldy**
- Spectral improvement: **Modified Foldy**



(n) Real 3D case with  $N_{\text{obs}} = 13$  (fixed distance)



(o) Aligned obstacles with  $N_{\text{obs}} = 5$  ( $\sqrt{\delta}$ -dependence)



(p) Aligned obstacles with  $N_{\text{obs}} = 5$  ( $\delta$ -dependence)

# Outline

## 1. Asymptotic models

- Single-scattering
- Application: Born approximation
- Multiple-scattering: Foldy-Lax model
- Numerical results

## 2. Spectral method: Spherical case

- Discretization
- Numerical convergence
- Comparison with asymptotic models
- Comparison with finite element solutions

## 3. Conclusions and perspectives



# Spectral method

- The electromagnetic fields are represented by the **Stratton-Chu formula** for  $\mathbf{x} \in \Omega_\delta$

$$\mathbf{E}_\delta(\mathbf{x}) = \sum_{k=1}^{N_{\text{obs}}} \mathbf{curl} \int_{\Gamma_\delta^k} \Phi(\mathbf{x}, \mathbf{y}) \mathbf{p}_k(\mathbf{y}) d\mathbf{s}_\mathbf{y}$$

where  $\Phi(\mathbf{x}, \mathbf{y}) = \frac{\exp(i\kappa|\mathbf{x}-\mathbf{y}|)}{4\pi|\mathbf{x}-\mathbf{y}|}$

# Spectral method

- The electromagnetic fields are represented by the **Stratton-Chu formula** for  $\mathbf{x} \in \Omega_\delta$

$$\mathbf{E}_\delta(\mathbf{x}) = \sum_{k=1}^{N_{\text{obs}}} \mathbf{curl} \int_{\Gamma_\delta^k} \Phi(\mathbf{x}, \mathbf{y}) \mathbf{p}_k(\mathbf{y}) d\mathbf{s}_\mathbf{y}$$

- Each  $\mathbf{p}_k \in \mathbf{H}_t^{-\frac{1}{2}}(\text{div}_{\Gamma_\delta^k}, \Gamma_\delta^k)$  satisfies the **boundary integral equation**

$$\sum_{j=1}^{N_{\text{obs}}} \mathcal{M}_\Gamma^{kj} \mathbf{p}_j = -\mathbf{n} \times \mathbf{E}^{\text{inc}} \quad \text{on } \Gamma_\delta^k$$

where  $\mathcal{M}_\Gamma^{kj} : \mathbf{H}_t^{-\frac{1}{2}}(\text{div}_{\Gamma_\delta^j}, \Gamma_\delta^j) \longrightarrow \mathbf{H}_t^{-\frac{1}{2}}(\text{div}_{\Gamma_\delta^k}, \Gamma_\delta^k)$  is the extension of

$$\mathcal{M}_\Gamma^{kj} \lambda(\mathbf{x}_\Gamma) = \mathbf{n}(\mathbf{x}_\Gamma) \times \lim_{\mathbf{x} \rightarrow \mathbf{x}_\Gamma} \mathbf{curl} \int_{\Gamma_\delta^j} \Phi(\mathbf{x}, \mathbf{y}) \lambda(\mathbf{y}) d\mathbf{s}_\mathbf{y} \quad \lambda \in \mathcal{C}^\infty(\Gamma_\delta^j) \quad \mathbf{x}_\Gamma \in \Gamma_\delta^k$$

# Spectral method

- The electromagnetic fields are represented by the **Stratton-Chu formula** for  $\mathbf{x} \in \Omega_\delta$

$$\mathbf{E}_\delta(\mathbf{x}) = \sum_{k=1}^{N_{\text{obs}}} \mathbf{curl} \int_{\Gamma_\delta^k} \Phi(\mathbf{x}, \mathbf{y}) \mathbf{p}_k(\mathbf{y}) d\mathbf{s}_\mathbf{y}$$

- Each  $\mathbf{p}_k \in \mathbf{H}_t^{-\frac{1}{2}}(\text{div}_{\Gamma_\delta^k}, \Gamma_\delta^k)$  satisfies the **boundary integral equation**

$$\sum_{j=1}^{N_{\text{obs}}} \mathcal{M}_{\Gamma}^{kj} \mathbf{p}_j = -\mathbf{n} \times \mathbf{E}^{\text{inc}} \quad \text{on } \Gamma_\delta^k$$

- Galerkin discretization** of the BIE on **local spectral basis** with  $N_{\text{mod}}$  modes

$$\mathbf{p}_j(\mathbf{x}) = \sum_{n=1}^{N_{\text{mod}}} \sum_{m=-n}^n p_{n,m}^{j,\perp} \nabla_{S^2} Y_{n,m}(\hat{\mathbf{x}}_j) + p_{n,m}^{j,\times} \mathbf{curl}_{S^2} Y_{n,m}(\hat{\mathbf{x}}_j) \quad \mathbf{x} \in \Gamma_\delta^j$$

with  $\hat{\mathbf{x}}_j = \frac{\mathbf{x} - \mathbf{c}_j}{|\mathbf{x} - \mathbf{c}_j|}$  and  $\nabla_{S^2} Y_{n,m}$ ,  $\mathbf{curl}_{S^2} Y_{n,m}$ : complex-valued vector spherical harmonics

# Vectorial formulation

- **Variational** formulation: Find  $\mathbf{p}_k \in \mathbf{H}_t^{-\frac{1}{2}}(\text{div}_{\Gamma_\delta^k}, \Gamma_\delta^k)$  such that

$$\sum_{j=1}^{N_{\text{obs}}} \langle \mathcal{M}_{\Gamma}^{kj} \mathbf{p}_j, \mathbf{v}_k \rangle_{\Gamma_\delta^k} = - \langle \mathbf{n} \times \mathbf{E}^{\text{inc}}, \mathbf{v}_k \rangle_{\Gamma_\delta^k} \quad \forall \mathbf{v}_k \in \mathbf{H}_t^{-\frac{1}{2}}(\text{curl}_{\Gamma_\delta^k}, \Gamma_\delta^k)$$

- **Vectorial** formulation: Find  $\mathbf{p} = ((p_{n,m}^{1,\perp}), \dots, (p_{n,m}^{N_{\text{obs}},\perp}), (p_{n,m}^{1,\times}), \dots, (p_{n,m}^{N_{\text{obs}},\times}))^\top \in \mathbb{C}^{\mathbf{N}}$  s.t.

$$\mathbb{M} \mathbf{p} = \mathbf{f}$$

with  $\mathbf{N} = 2N_{\text{mod}}(N_{\text{mod}} + 2)N_{\text{obs}}$  and

$$\mathbb{M} = \begin{pmatrix} \mathbb{M}_{\perp\perp} & \mathbb{M}_{\perp\times} \\ \mathbb{M}_{\times\perp} & \mathbb{M}_{\times\times} \end{pmatrix} \quad \text{with} \quad \mathbb{M}_{\alpha\beta} = \begin{pmatrix} \mathbb{M}_{\alpha\beta}^{11} & \mathbb{M}_{\alpha\beta}^{12} & \dots & \mathbb{M}_{\alpha\beta}^{1N_{\text{obs}}} \\ \mathbb{M}_{\alpha\beta}^{21} & \mathbb{M}_{\alpha\beta}^{22} & \dots & \mathbb{M}_{\alpha\beta}^{2N_{\text{obs}}} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbb{M}_{\alpha\beta}^{N_{\text{obs}}1} & \mathbb{M}_{\alpha\beta}^{N_{\text{obs}}2} & \dots & \mathbb{M}_{\alpha\beta}^{N_{\text{obs}}N_{\text{obs}}} \end{pmatrix}$$

$$= \mathbb{M}_{\alpha\beta}(\delta, \mathbf{d}_{jk})$$

# Vectorial formulation

- Variational formulation: Find  $\mathbf{p}_k \in \mathbf{H}_t^{-\frac{1}{2}}(\text{div}_{\Gamma_\delta^k}, \Gamma_\delta^k)$  such that

$$\sum_{j=1}^{N_{\text{obs}}} \langle \mathcal{M}_{\Gamma}^{kj} \mathbf{p}_j, \mathbf{v}_k \rangle_{\Gamma_\delta^k} = -\langle \mathbf{n} \times \mathbf{E}^{\text{inc}}, \mathbf{v}_k \rangle_{\Gamma_\delta^k} \quad \forall \mathbf{v}_k \in \mathbf{H}_t^{-\frac{1}{2}}(\text{curl}_{\Gamma_\delta^k}, \Gamma_\delta^k)$$

- Vectorial formulation: Find  $\mathbf{p} = ((p_{n,m}^{1,\perp}), \dots, (p_{n,m}^{N_{\text{obs}},\perp}), (p_{n,m}^{1,\times}), \dots, (p_{n,m}^{N_{\text{obs}},\times}))^\top \in \mathbb{C}^N$  s.t.

$$\mathbb{M} \mathbf{p} = \mathbf{f}$$



Computation time  
Ill-conditioned system  
Memory consumption

- Numerical integration (Gauss-Lobatto)
- Large number of unknowns
- Dense matrix



Mex interface (C++)  
Preconditioning (system & matrix)  
Smart storage and assembling

- Code speed-up
- Linear algebra tools
- Sub-blocks  $\mathbb{M}_{\alpha\beta}^{kj}$  depend on  $\mathbf{d}_{jk}$

# Preconditionning

Linear system: **Change of variable** + **Permutation** + **Normalization**

$$\mathbf{E}_\delta = \phi_\delta(\mathbf{p}_\delta) \quad \text{with } \mathbb{M} \mathbf{p}_\delta = \mathbf{f}$$



$$\mathbf{E}_\delta = \phi(\tilde{\mathbf{p}}_\delta) \quad \text{with } \tilde{\mathbb{M}} \tilde{\mathbf{p}}_\delta = \tilde{\mathbf{f}}$$



Dense matrix  $\tilde{\mathbb{M}}$



Error of approximation

Analytic preconditionner (e.g. dipole)

Algebraic preconditionner (small criterion)

# Preconditionning

Linear system: Change of variable + Permutation + Normalization

$$\mathbf{E}_\delta = \phi_\delta(\mathbf{p}_\delta) \quad \text{with } \mathbb{M} \mathbf{p}_\delta = \mathbf{f}$$



$$\mathbf{E}_\delta = \phi(\tilde{\mathbf{p}}_\delta) \quad \text{with } \tilde{\mathbb{M}} \tilde{\mathbf{p}}_\delta = \tilde{\mathbf{f}}$$




Dense matrix  $\tilde{\mathbb{M}}$




Error of approximation

Analytic preconditionner (e.g. dipole)


Algebraic preconditionner (small criterion)



(a)  $N_{\text{obs}} = 4$  and  $N_{\text{mod}} = 1$



(b)  $N_{\text{obs}} = 25$  and  $N_{\text{mod}} = 1$



(c)  $N_{\text{obs}} = 4$  and  $N_{\text{mod}} = 5$

Figure: Skeleton of  $\tilde{\mathbb{M}}$

# Preconditionning

Linear system: Change of variable + Permutation + Normalization

$$\mathbf{E}_\delta = \phi_\delta(\mathbf{p}_\delta) \quad \text{with } \mathbb{M} \mathbf{p}_\delta = \mathbf{f}$$



$$\mathbf{E}_\delta = \phi(\tilde{\mathbf{p}}_\delta) \quad \text{with } \tilde{\mathbb{M}} \tilde{\mathbf{p}}_\delta = \tilde{\mathbf{f}}$$




Dense matrix  $\tilde{\mathbb{M}}$




Error of approximation

Analytic preconditionner (e.g. dipole)


Algebraic preconditionner (small criterion)



(a)  $N_{\text{obs}} = 4$  and  $N_{\text{mod}} = 1$



(b)  $N_{\text{obs}} = 25$  and  $N_{\text{mod}} = 1$



(c)  $N_{\text{obs}} = 4$  and  $N_{\text{mod}} = 5$

Figure: Skeleton of  $\tilde{\mathbb{M}}$



# Preconditionning

Linear system: Change of variable + Permutation + Normalization

$$\mathbf{E}_\delta = \phi_\delta(\mathbf{p}_\delta) \quad \text{with } \mathbb{M} \mathbf{p}_\delta = \mathbf{f}$$



$$\mathbf{E}_\delta = \phi(\tilde{\mathbf{p}}_\delta) \quad \text{with } \tilde{\mathbb{M}} \tilde{\mathbf{p}}_\delta = \tilde{\mathbf{f}}$$




Dense matrix  $\tilde{\mathbb{M}}$




Error of approximation

Analytic preconditionner (e.g. dipole)


Algebraic preconditionner (small criterion)



(a)  $N_{\text{obs}} = 4$  and  $N_{\text{mod}} = 1$



(b)  $N_{\text{obs}} = 25$  and  $N_{\text{mod}} = 1$



(c)  $N_{\text{obs}} = 4$  and  $N_{\text{mod}} = 5$

Figure: Skeleton of  $\text{Precond}(\tilde{\mathbb{M}})$

# Preconditionning

Linear system: Change of variable + Permutation + Normalization

$$\mathbf{E}_\delta = \phi_\delta(\mathbf{p}_\delta) \quad \text{with } \mathbb{M} \mathbf{p}_\delta = \mathbf{f}$$



$$\mathbf{E}_\delta = \phi(\tilde{\mathbf{p}}_\delta) \quad \text{with } \tilde{\mathbb{M}} \tilde{\mathbf{p}}_\delta = \tilde{\mathbf{f}}$$




Dense matrix  $\tilde{\mathbb{M}}$




Error of approximation

Analytic preconditionner (e.g. dipole)


Algebraic preconditionner (small criterion)



(a)  $N_{\text{obs}} = 4$  and  $N_{\text{mod}} = 1$



(b)  $N_{\text{obs}} = 25$  and  $N_{\text{mod}} = 1$



(c)  $N_{\text{obs}} = 4$  and  $N_{\text{mod}} = 5$


Figure: Skeleton of  $\text{Precond}(\tilde{\mathbb{M}})$

# Smart storage and assembling

- Example: 4 aligned obstacles




$$\tilde{\mathbf{M}} =$$



The sub-blocks  $\mathbf{M}_{\alpha\beta}^{kj}$  depend only on  $\delta$  and  $\mathbf{d}_{jk}$

## Smart storage and assembling

- Example: 4 aligned obstacles



$$\tilde{\mathbf{M}} =$$

Figure 1 shows four 4x4 grids illustrating the evolution of a quantum state. Each grid has a diagonal of dots from top-left to bottom-right. The grids are labeled as follows:


- Grid 1 (top-left): 'Id' at (1,1) and (4,4), and '1 - 2' at (1,2).
- Grid 2 (top-right): '0' at (1,1) and (4,4), and '1' at (1,2), (2,3), and (3,4).
- Grid 3 (bottom-left): '0' at (1,1) and (4,4), and '1' at (1,2), (2,3), and (3,4).
- Grid 4 (bottom-right): 'Id' at (1,1) and (4,4), and '1' at (1,2), (2,3), and (3,4).

- Storage of sub-blocks

[illegible]

## Smart storage and assembling

- Example: 4 aligned obstacles



$$\tilde{\mathbf{M}} =$$

Id	1 - 2	1 - 3	
			Id

0			
			0

0			
			0

Id			
			Id


- Storage of sub-blocks

[illegible]



## Smart storage and assembling

- Example: 4 aligned obstacles




$$\tilde{\mathbf{M}} =$$

- Storage of sub-blocks

[illegible]

## Smart storage and assembling

- Example: 4 aligned obstacles



$$\tilde{\mathbf{M}} =$$

Id	1 - 2	1 - 3	1 - 4
2 - 1	.	.	.
3 - 1	.	.	.
	.	.	Id

0	.	.	.
.	.	.	.
.	.	.	.
	.	.	0


- Storage of sub-blocks

[illegible]



## Smart storage and assembling

- Example: 4 aligned obstacles




$$\tilde{\mathbf{M}} =$$

- Storage of sub-blocks

[illegible]

## Smart storage and assembling

- Example: 4 aligned obstacles



$$\tilde{M} =$$

Figure 1: A 4x4 grid of 16 cells, each containing a 4x4 grid of 16 smaller cells. The top-left cell of the large grid is labeled 'Id' and contains a 4x4 grid of colored squares (orange, light orange, light orange, red) with labels '1 - 2', '1 - 3', and '1 - 4'. The top-right cell is labeled '0' and contains a 4x4 grid of colored squares (light green, light green, light green, dark green). The bottom-left cell is labeled '0' and contains a 4x4 grid of colored squares (light blue, light blue, light blue, dark blue). The bottom-right cell is labeled 'Id' and contains a 4x4 grid of colored squares (pink, pink, pink, dark pink). The middle two rows of the large grid contain 4x4 grids of colored squares with labels '2 - 1', '3 - 1', and '4 - 1' in the top-left cell of each row.

- Storage of sub-blocks

[illegible]




- Iterative solvers: define the **action** of  $\tilde{\mathbf{M}}_{\text{Block}}$  on  $\mathbf{u}$

$$\mathcal{A}(\tilde{\mathbb{M}}_{\text{Block}}, \mathbf{u}) := \tilde{\mathbb{M}} \mathbf{u}$$





# Numerical validation of spectral solutions

- Incident field: Electromagnetic plane wave along z-axis polarized by  $\mathbf{e}_x$
- Reference solution: Spectral solution with  $N_{\text{mod}} = 10$



# Comparison with asymptotic models




Difference between Spectral 1 and modified Foldy



	Modified Foldy				Spectral 1			
$N_{\text{obs}}$	100	1000	5000	10000	100	1000	5000	10000
<b>Linear system</b>	0.17	17.71	34.01	43.49	2.66	40.74	352.47	--
<b>Post-processing</b>	5.25	47.84	254.34	459.90	12.17	126.81	> 3600	--

Table: Time comparison in seconds

# Comparison with finite element solutions (Montjoie)



Spectral 3			
$N_{\text{obs}}$	1	2	4
Size of linear system	30	60	120
$L^2$ -Accuracy ( $\times 10^{-6}$ )	4.77	4.48	3.66
<b>Time</b> (s)	15.46	65.55	262.33
<b>Memory</b> (GB)	0.3	2.36	9.3

Finite Element 3			
$N_{\text{obs}}$	1	2	4
Size of linear system	925 422	1 972 800	3 942 936
$L^2$ -Accuracy ( $\times 10^{-3}$ )	0.28	2.27	3.31
<b>Time</b> (min)	26.5	54.25	147.25
<b>Memory</b> (GB)	21.11	36.97	75

Table: Computational costs

# Outline

## 1. Asymptotic models

- Single-scattering
- Application: Born approximation
- Multiple-scattering: Foldy-Lax model
- Numerical results

## 2. Spectral method: Spherical case

- Discretization
- Numerical convergence
- Comparison with asymptotic models
- Comparison with finite element solutions

## 3. Conclusions and perspectives

# Conclusions and Perspectives

## Conclusion

- ✓ Derivation and numerical validation of the matched asymptotic expansions
- ✓ Extension to Born and Foldy-Lax models
- ✓ Derivation and implementation at any order spectral method + Numerical validation

## On-going work

- ~ Comparison of preconditionners and iterative solvers
- ~ Smart assembling for particular configurations (plane, cubic volume)
- ~ Contribution to mathematical justification of the asymptotic expansions

## Perspectives

- ✗ Existence of electromagnetic centers?
- ✗ Extension to obstacles of arbitrary shape
- ✗ Extension to time-dependent domain



Thank you for your attention